






# Scuttle flies (Diptera, Phoridae) collected by mosquito trap from Košice Zoo, Central Europe

Bernd Grundmann<sup>1</sup>, Katarína Loziaková Peňazziová<sup>2</sup>, Tomáš Csank<sup>2</sup>, Laura Mlynárová<sup>3</sup>, Patrik Pastorek<sup>4</sup>, Jozef Oboňa<sup>3</sup>

<sup>1</sup> Diekstraße 6, D – 33824 Werther (Westf.), Germany

<sup>2</sup> Department of Microbiology and Immunology, University of Veterinary Medicine and Pharmacy, Komenského 73, SK – 04181 Košice, Slovakia

<sup>3</sup> Department of Ecology, Faculty of Humanities and Natural Sciences, University of Prešov, 17. novembra 1, SK – 08116 Prešov, Slovakia

<sup>4</sup> ZOO Košice, Ulica k Zoologickej Záhrade 1, SK – 04001 Košice, Slovakia

Corresponding author: Laura Mlynárová ([laura.mlynarova@smail.unipo.sk](mailto:laura.mlynarova@smail.unipo.sk), [l.mlynarova93@gmail.com](mailto:l.mlynarova93@gmail.com))

## Abstract

In this paper the results of a Phoridae collection from Košice Zoo in eastern Slovakia are reported. An unusual trap method was employed – BG-Sentinel 2 mosquito traps combined with CO<sub>2</sub> as an attractant. Between July and October 2023, 73 species of the Phoridae family were recorded, 24 of which are new to the Slovak fauna (1 species from the genus *Woodiphora* and 23 species from the genus *Megaselia*). As a result, the total number of Phoridae species documented in Slovakia has increased to 254.

**Key words:** Faunistic, mosquito BG-Sentinel 2 trap, new records, Phoridae, Slovakia



Academic editor: Torsten Dikow

Received: 15 January 2025

Accepted: 6 May 2025

Published: 4 July 2025

ZooBank: <https://zoobank.org/A79B32A5-F39A-40AE-A495-0663485C80E6>

Citation: Grundmann B, Loziaková K, Peňazziová K, Csank T, Mlynárová L, Pastorek P, Oboňa J (2025) Scuttle flies (Diptera, Phoridae) collected by mosquito trap from Košice Zoo, Central Europe. ZooKeys 1244: 41–60. <https://doi.org/10.3897/zookeys.1244.146861>

Copyright: © Bernd Grundmann et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

Phorid flies (Diptera: Phoridae) belong to the suborder Cyclorrhapha, a superfamily within Platypteroidea, comprising approximately 35 genera and more than 700 species in Europe (Oosterbroek, 2006). To date, 230 species have been recorded from Slovakia (Mocek 1997, 2009; Jančík and Disney 2020; Grundmann et al. 2023, 2024). These flies, commonly known as scuttle flies, are characterised by their rapid, somewhat abrupt movements. Adult flies are typically small to medium-sized (0.5–6 mm) and often exhibit a somewhat curved, robust body form (Disney 1983). Most adults feed on nectar, honeydew, sap from fresh carrion, and faeces; some also feed on the body fluids of live beetle larvae and pupae.

According to Disney (1994), Phoridae can be collected by using a wide variety of insect trapping techniques. A commonly employed method is the use of Malaise traps (e.g. Grundmann and Kappert 2023; Caruso et al. 2024). Other methods, such as emergence traps (Szadziawska 1977; Büchs 1988), white and yellow water traps, pitfall traps (Disney et al. 1981), and light traps (Brown and Marshall 1984) are also frequently used. Breeding scuttle flies from natural substrates such as humus, compost, mushrooms, and carrion is another effective method (Disney and Evans 1982; Disney 1994). However, to the authors' knowledge, mosquito traps have not yet been used to capture scuttle flies.



## Materials and methods

### Sampling

A fatal case of West Nile virus infection in a Great Grey Owl (*Strix nebulosi* Forster, 1772) kept at Košice Zoo was diagnosed (Peňazziová et al. 2021). Based on this, mosquito capture was conducted using BG-Sentinel 2 traps (Biogents, Germany) with CO<sub>2</sub> cylinders as an attractant (Fig. 1). The traps were placed near small lakes in the locality (see locality data) and operated continuously from July 2023 (with the first collection on 17 July) until the end of October 2023 (with last collection on 28 October). The capture nets were replaced twice a week and stored at -20 °C until transport to the laboratory, where they were stored to -80 °C. After sorting the mosquitoes, the remaining material was preserved in 75% alcohol. The collected flies were sorted to family level and identified to species level by BG using identification keys (Schmitz, 1941, 1943, 1951, 1956; Disney, 1983, 1989, 1994, 1999), who also curated the collection.

Locality data:

Slovakia, Košice district, Košice Zoo,

Horný rybník – Upper pond (HR): 48°47'11.4"N 21°12'11.9"E, 412 m a.s.l.

Dolný rybník – Lower pond (DR): 48°47'20.6"N 21°12'22.0"E, 411 m a.s.l.

An asterisk (\*) before the species name indicates a new geographical record for Slovakia. References to the online GBIF page resources (with a brief comment on their European distribution) are added for each species. For species new to the Slovak fauna, additional information on their known distribution and the feeding ecology of their larvae (if available) is also included.



**Figure 1.** Photograph of BG-Sentinel 2 trap (Biogents, Germany) from Košice Zoo.



## Results

A total of 487 scuttle flies were collected in the BG-Sentinel 2 traps (July – 100 individuals, August – 96 inds, September – 180 inds, October – 111 inds) in 2023, representing 73 species (see Annotated list of recorded species). Thirty individuals were indeterminate females of the genus *Megaselia* and six individuals belonged to three new species. Twenty-four species were recorded for the first time in Slovakia, and six species represent the second record for Slovakian fauna. The most abundant species were *Megaselia angusta* (Wood, 1909) with 69 individuals, *Megaselia frameata* Schmitz, 1927 with 65 inds, and *Megaselia aculeata* (Schmitz, 1919) with 41 inds. Twenty-seven species were recorded based on only one individual.

### Annotated list of recorded species

#### Diptera

#### Family: Phoridae

##### *Anevrina thoracica* (Meigen, 1804)

**Material examined.** DR 09.10. ♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only rarely in Central Europe. <https://www.gbif.org/species/1545916>.

##### *Conicera dauci* (Meigen, 1830)

**Material examined.** DR 23.10. ♂; HR 22.08. ♂, 28.08. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1546041>.

##### *Conicera floricola* Schmitz, 1938

**Material examined.** DR 31.07. ♂, 28.08. ♂, 11.09. ♂, 26.09. ♂; HR 04.09. ♂, 07.09. ♂, 22.09. ♂, 03.10. ♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only sporadically in Central Europe. <https://www.gbif.org/species/1546039>.

##### *Conicera tibialis* Schmitz, 1925

**Material examined.** DR 04.09. 2 ♂♂, 07.09. 2 ♂♂, ♀, 11.09. ♂; HR 31.07. ♂, 11.09. ♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions. <https://www.gbif.org/species/1546012>.

##### *Diplonevra nitidula* (Meigen, 1830)

**Material examined.** DR 03.08. ♀, 09.08. ♂, 11.08. ♂, 04.09. 2 ♂♂, ♀, 13.10. ♂, ♀.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only sporadically in Central Europe. <https://www.gbif.org/species/1548598>.



***Gymnophora arcuata* (Meigen, 1830)**

**Material examined.** DR 31.07. ♂, 09.10. ♂; HR 22.08. ♀.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only sporadically in Central Europe. <https://www.gbif.org/species/1549517>.

***Pseudacteon formicarum* (Verrall, 1877)**

**Material examined.** DR 17.07. ♂, 26.07. ♂.

**GBIF records.** In Europe this species is rarely recorded only the northern regions. <https://www.gbif.org/species/1549197>.

***Triphleba antricola* (Schmitz, 1918)**

**Material examined.** HR 18.09. ♀.

**GBIF records.** In Europe this species is sporadically recorded in Central Europa and in the northwest of the Mediterranean region. <https://www.gbif.org/species/1549712>.

***Triphleba distinguenda* (Strobl, 1892)**

**Material examined.** HR 22.08. ♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only sporadically in Central Europe. <https://www.gbif.org/species/1549634>.

***Triphleba dudai* (Schmitz, 1918)**

**Material examined.** HR 28.08. ♂, 03.10. ♂.

**GBIF records.** There are only a few records from Sweden. <https://www.gbif.org/species/1549630>.

***Triphleba lugubris* (Meigen, 1830)**

**Material examined.** DR 09.10. 3 ♂♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions. <https://www.gbif.org/species/1549637>.

***Triphleba nudipalpis* (Becker, 1901)**

**Material examined.** HR 21.07. ♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions and only sporadically in Central Europe. <https://www.gbif.org/species/9748380>.

**\**Woodiphora retroversa* (Wood, 1908)**

**Material examined.** HR 26.07. ♂.

**Published records.** Distribution in Britain, Denmark (mainland), Hungary, and the Netherlands, with records from Poland (Durska 2013).



**GBIF records.** Additional distribution in Norway. <https://www.gbif.org/species/1546062>.

**Comments.** This is a summer species, most abundant in July. Its development occurs in carrion (Schmitz 1956).

***Megaselia aculeata* (Schmitz, 1919)**

**Material examined.** DR 17.07. ♂, 18.08. 2 ♂♂, 04.09. 2 ♂♂, 07.09. ♂, 11.09. ♂, 18.09. 3 ♂♂, 26.09. ♂, 09.10. 3 ♂♂, 10.10. ♂, 13.10. 2 ♂♂, 23.10. 3 ♂♂; HR 22.08. ♂, 11.09. ♂, 14.09. ♂, 22.09. ♂, 03.10. 10 ♂♂, 13.10. 3 ♂♂, 28.10. 4 ♂♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547024>.

**Comments.** The first Slovak record of this species was published by Grundmann et al. (2024).

***Megaselia aequalis* (Wood, 1909)**

**Material examined.** DR 14.09. ♀; HR 11.09. 2 ♂♂.

**GBIF records.** In Europe this species is mainly recorded in the northern regions. <https://www.gbif.org/species/1546813>.

**\**Megaselia albicans* (Wood, 1908)**

**Material examined.** HR 28.07. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe and the Balkans. <https://www.gbif.org/species/1547725>.

**Comments.** The larvae of this species are reported to be mycophagous (Durska 2013). It has been reared from false morel (*Gyromitra esculenta* (Pers.) Fr.) a mushroom of the family Morchellaceae (Disney 1994).

***Megaselia albocingulata* (Strobl, 1906)**

**Material examined.** HR 17.07. 2 ♂♂, 18.08. ♂; DR 17.07. ♂, 31.07. ♂, 11.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1547320>.

**\**Megaselia analis* (Lundbeck, 1920)**

**Material examined.** DR 26.09. ♂.

**GBIF records.** There are only a few records from Finland and Sweden. <https://www.gbif.org/species/1548247>.

**Comments.** The larval diet is unknown (Durska 2013).

***Megaselia angularis* (Schmitz, 1924)**

**Material examined.** DR 14.08. ♂, 18.08. 2 ♂♂; HR 17.07. ♂, 18.08. ♂, 22.08. 2 ♂♂.



**GBIF records.** There are only a few records from Finland and Sweden. <https://www.gbif.org/species/1546855>.

***Megaselia angusta* (Wood, 1909)**

**Material examined.** DR 28.07. ♂, 09.08. ♂, 28.08. ♂, 04.09. ♀, 11.09. 2 ♂, 14.09. ♂, 18.09. 4 ♂♂, 22.09. 2 ♂♂, 09.10. ♂, 10.10. ♂, 28.10. 2 ♂♂; HR 17.07. ♂, 21.07. 3 ♂♂, 24.07. 12 ♂♂, 28.07. ♂, 07.08. 3 ♂♂, 3 ♀♀, 25.08. ♂, 28.08. 2 ♂♂, 04.09. 3 ♂♂, 07.09. 16 ♂♂, 11.09. ♂, 14.09. ♂, 22.09. ♂, 26.09. ♂, 03.10. 2 ♂♂, 28.10. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1547218>.

**\**Megaselia annulipes* (Schmitz, 1921)**

Fig. 2

**Material examined.** DR 14.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe and the Balkans. <https://www.gbif.org/species/1546937>.

**Comments.** The larva is a parasitoid of spiders. It has been reared from *Moebelia penicillata* (Westring, 1851), a spider of the family Linyphiidae (Weber et al. 2006).

**\**Megaselia basispinata* (Lundbeck, 1920)**

**Material examined.** DR 09.08. ♂.

**Published records.** This Holarctic-Neotropical species is known from Europe, Macaronesia, and North and South America (Langourov 2004a).

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1548403>.

**Comments.** The larval diet is unknown (Durska 2013).

***Megaselia berndseni* (Schmitz, 1919)**

**Material examined.** HR 26.07. 2 ♂♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/4516507>.

***Megaselia brevicostalis* (Wood, 1910)**


**Material examined.** DR 18.08. ♂; HR 21.07. ♂, 22.08. ♂, 28.08. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547644>.

**\**Megaselia brevissima* Schmitz, 1924**

**Material examined.** DR 31.07. ♂.





**Figure 2.** *Megaselia annulipes* (Schmitz, 1921). This species is easy recognisable by its annulated front tarsi, although little is known about their function. A signalling function in the context of mating behaviour could be conceivable (photograph by [www.spessart-fliegen.de](http://www.spessart-fliegen.de)). Scale bar 1 mm.

**GBIF records.** No distribution records are currently shown on the map. <https://www.gbif.org/species/1547213>.

**Comments.** Identification with Disney (2006a). This species belongs to the *Megaselia breviterga* species complex.

***Megaselia breviterga* (Lundbeck, 1920)**

**Material examined.** DR 26.07. ♀

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1548369>.

**Comments.** The first record was published by Grundmann et al. (2023).

***Megaselia ciliata* (Zetterstedt, 1848)**

**Material examined.** HR 25.08. ♂; 07.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern and Central European regions. <https://www.gbif.org/species/1546686>.

***Megaselia cinereifrons* (Strobl, 1910)**

**Material examined.** HR 21.07. 2 ♂♂; HR 31.07. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1548194>.



***Megaselia communiformis* (Schmitz, 1918)**

**Material examined.** HR 07.08. ♂.

**GBIF records.** There are only few records from Sweden and Germany. <https://www.gbif.org/species/1547173>.

***Megaselia conformis* (Wood, 1909)**

**Material examined.** DR 07.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1548180>.

**\**Megaselia crassipes* (Wood, 1909)**

**Material examined.** HR 28.08. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1546701>.

**Comments.** The larval diet is unknown (Durska 2013).

**\**Megaselia curvicapilla* Schmitz, 1947**

**Material examined.** HR 28.08. ♂, HR 04.09. ♂.

**GBIF records.** There are only few records from Finland, Sweden, and Germany. <https://www.gbif.org/species/1546685>.

**Comments.** The larval diet is unknown (Durska 2013).

**\**Megaselia deltomera* (Schmitz, 1924)**

**Material examined.** HR 21.07. ♂, HR 22.09. ♂.

**GBIF records.** No distribution records are currently shown on the map. <https://www.gbif.org/species/1547749>.

**Comments.** Identification with Disney (2004).

**\**Megaselia densior* Schmitz, 1927**

**Material examined.** DR 31.07. ♂, 11.09. 2 ♂♂, 22.09. 2 ♂♂, 26.09. ♂, 09.10. ♂; HR 26.07. ♂; 28.07. ♂, 07.08. ♂, 22.08. ♂, 11.09. 2 ♂♂, 14.09. ♂.

**GBIF records.** No distribution records are currently shown on the map. <https://www.gbif.org/species/1547247>.

**Comments.** This is the eponymous species of the *Megaselia densior* species complex, which has been revised by Buck and Disney (2001).

***Megaselia emarginata* (Wood, 1908)**

**Material examined.** DR 21.07. ♂, DR 26.07. ♂; HR 25.08. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1546757>.



***Megaselia errata* (Wood, 1912)**

**Material examined.** DR 28.07. ♂, DR 18.08. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1548095>.

**\**Megaselia excorticata* Disney, 2010**

**Material examined.** DR 11.09. ♂, 10.10. ♂; HR 26.07. ♂, 07.08. ♂, 18.08. ♂, 13.10. ♂.

**Published records.** In addition to the type locality in Finland, there is only one further record from Germany (Hannig et al. 2023).

**GBIF records.** There are only few records from Finland. <https://www.gbif.org/species/11161573>.

**Comments.** Identification with Disney (2010a).

**\**Megaselia fenestralis* (Schmitz, 1919)**

**Material examined.** DR 21.07. ♂, 14.08. ♂, 25.08. ♂, 28.08. ♂; HR 17.07. ♂.

**GBIF records.** There are only few records from Finland and Sweden. <https://www.gbif.org/species/1547966>.

**Comments.** The larval diet is unknown (Durska 2013).

***Megaselia flava* (Fallén, 1823)**

**Material examined.** HR 18.09. ♀.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547612>.

**\**Megaselia flavescens* (Wood, 1909)**

**Material examined.** DR 21.07. ♂.

**GBIF records.** There are only few records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1547805>.

**Comments.** The larval diet is unknown (Durska 2020).

***Megaselia flavicans* Schmitz, 1935**

**Material examined.** DR 31.07. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1546504>.

***Megaselia frameata* Schmitz, 1927**

**Material examined.** DR 26.09. ♂, 09.10. 2 ♂♂, 23.10. 2 ♂; HR 21.07. ♂, 24.07. 2 ♂, 22.08. ♂, 25.08. 2 ♂, 04.09. ♂, 04.09. ♂, 14.09. ♂, 18.09. 3 ♂♂, 22.09. 3 ♂♂, 26.09. 17 ♂♂, 03.10. 22 ♂♂, 13.10. ♂, 28.10. 3 ♂♂.



**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547462>.

***Megaselia giraudii* (Egger, 1862)**

**Material examined.** DR 25.08. ♂, 28.08. ♂, 11.09. ♂, 14.09. ♂, 09.10. 2 ♂♂; HR 17.07. 2 ♂♂, 21.07. ♂, 24.07. ♂, 18.08. 2 ♂♂, 22.08. ♂, 25.08. 2 ♂, 28.08. 2 ♂, 04.09. ♂, 07.09. ♂, 11.09. 5 ♂♂, 26.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/4295479>.

***Megaselia gregaria* (Wood, 1910)**

**Material examined.** DR 17.07. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1546523>.

**\**Megaselia hartfordensis* Disney, 1983**

**Material examined.** DR 21.07. ♂; HR 14.09. ♂.

**Published records.** In addition to the type locality in England, there is one record from France (Disney and Withers 2011) and one from Germany (Hannig et al. 2023).

**GBIF records.** There is only one record from Britain. <https://www.gbif.org/species/1546633>.

***Megaselia hilaris* Schmitz, 1927**

**Material examined.** HR 28.08. ♂.

**GBIF records.** In Europe this species is recorded in the northern and central regions. <https://www.gbif.org/species/1546935>.

**\**Megaselia ignobilis* (Schmitz, 1919)**

**Material examined.** DR 17.07. 2 ♂♂, 03.08. ♂, 25.08. ♂, 07.09. ♂, 11.09. ♂, 09.10. ♂; HR 24.07. 2 ♂♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1546714>.

**Comments.** The larval diet is unknown (Durska 2015, 2020).

**\**Megaselia intergeriva* Schmitz, 1948**

**Material examined.** HR 28.08. ♂.

**Published records.** Identification followed Disney (2010b). This species has been described from Austria. Disney (2010b) mentions finds from Switzerland where flies have been reared from spruce trunks. It is also known from Germany (Hannig et al. 2023).



**GBIF records.** No distribution records are currently shown on the map. <https://www.gbif.org/species/1548170>.

***Megaselia involuta* (Wood, 1910)**

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe and the Balkans. <https://www.gbif.org/species/1548490>.

**Material examined.** HR 26.07. ♂, 28.07. ♂.

**\**Megaselia latipalpis* (Schmitz, 1921)**

**Material examined.** DR 26.07. ♂, 31.07. ♂, 07.09. 2 ♂♂, 11.09. ♂, 14.09. ♂; HR 24.07. ♂, 31.07. ♂, 25.08. ♂, 04.09. ♂, 07.09. 3 ♂, 11.09. ♂.

**Published records.** European species, known from Austria, Balkan Peninsula, Germany, Great Britain, and France (Langourov 2004b).

**GBIF records.** There are only few records from Sweden and Britain. <https://www.gbif.org/species/1546846>.

***Megaselia ledhuriensis* Brues, 1915**

*Megaselia subfuscipes* Schmitz, 1935. Syn.

**Material examined.** HR 03.10. ♂.

**GBIF records.** There are only few records Britain. <https://www.gbif.org/species/10139024>.

**Comments.** The first Slovak record of this species was published by Grundmann et al. (2024).

***Megaselia major* (Wood, 1912)**

**Material examined.** DR 31.07. ♂.

**GBIF records.** There are only a few records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1547851>.

**\**Megaselia malhamensis* Disney, 1986**

**Material examined.** DR 14.09. ♂; HR 17.07. ♂, 22.08. ♂.

**Published records.** This species has been described from England and is also known from Germany.

**GBIF records.** There are only few records from Finland, Sweden, Germany, and the Balkans. <https://www.gbif.org/species/1547856>.

**Comments.** Identification with Buck and Disney (2001).

***Megaselia manicata* (Wood, 1910)**

**Material examined.** DR 11.09. ♂; HR 31.07. ♂, 22.08. ♂, 25.08. ♂, 28.08. 2 ♂♂.



**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547792>.

**Comments.** The first Slovak record of this species was published by Grundmann et al. (2024).

***Megaselia maura* (Wood, 1910)**

**Material examined.** DR 13.10. ♂.

**GBIF records.** There are only a few records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1546902>.

***Megaselia nasoni* (Malloch, 1914)**

**Material examined.** HR 22.08. ♂.

**GBIF records.** There are several records from Finland, Sweden, and Britain. <https://www.gbif.org/species/4516511>.

***Megaselia nigra* (Meigen, 1830)**

**Material examined.** HR 26.07. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1547368>.

***Megaselia pectoralis* (Wood, 1910)**

**Material examined.** DR 13.10. ♂; HR 14.09. ♂, 26.09. ♂.

**GBIF records.** There are several records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1546740>.

***Megaselia pleuralis* (Wood, 1909)**

**Material examined.** DR 04.09. ♂, 18.09. 2 ♂, 09.10. ♂, 10.10. ♂; HR 17.07. ♂, 31.07. ♂, 18.09. ♂, HR 03.10. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1548494>.

***Megaselia propinqua* (Wood, 1909)**

**Material examined.** HR 21.07. ♂.

**GBIF records.** There are several records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1546995>.

***Megaselia pusilla* (Meigen, 1830)**

**Material examined.** DR 21.07. 2 ♂♂, 25.08. ♂, 07.09. 2 ♂♂, 14.09. ♂, 09.10. ♂, 10.10. ♂, 23.10. ♂; HR 22.08. ♂, 28.08. 2 ♂, 04.09. ♂, 09.10. ♂.



**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1548452>.

***Megaselia pygmaea* (Zetterstedt, 1848)**

**Material examined.** DR 03.08. ♂, 09.08. ♂, 11.09. ♂.

**GBIF records.** In Europe, this species is primarily recorded in the northern regions. <https://www.gbif.org/species/1547970>.

***Megaselia quadriseta* (Schmitz, 1918)**

**Material examined.** DR 28.07. ♂, 03.08. 2 ♂, 25.08. ♂, 07.09. ♂, 11.09. 3 ♂, 10.10. ♂; HR 21.07. ♂, 14.09. 3 ♂♂, 03.10. ♂, 28.10. 2 ♂♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with two records also from Central Europe. <https://www.gbif.org/species/1547785>.

***Megaselia rubescens* (Wood, 1912)**

**Material examined.** DR 09.10. ♂; HR 18.08. ♂, 11.09. ♂.

**GBIF records.** There are several records from Sweden and Britain. <https://www.gbif.org/species/1547312>.

**Comments.** The first Slovak record of this species was published by Grundmann et al. (2024).

***Megaselia rufipes* (Meigen, 1804)**

**Material examined.** DR 11.09. 2 ♂♂, 14.09. ♂.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe and the Balkans. <https://www.gbif.org/species/1547156>.

**\**Megaselia shawi* Disney, 2006**

**Material examined.** DR 09.10. ♂; HR 28.07. ♂, 04.09. ♂.

**Published records.** Identification with Disney (2006b). This species is a sibling species of the common *Megaselia frameata*. It is known only from the type locality in Scotland. Meanwhile, *M. shawi* proved to be an overlooked species, which is widely distributed in Europe (BG unpublished). The type specimens have been reared from a dead *Salix* trunk.

**GBIF records.** No distribution records are currently shown on the map. <https://www.gbif.org/species/1547691>.

**\**Megaselia speiseri* Schmitz, 1929**

**Material examined.** DR 26.09. ♂, 10.10. ♂.

**Published records.** This species has been recorded as new for the German fauna by Prescher and Bellstedt (1994).



**GBIF records.** There are several records from Sweden, Finland, Denmark, and Germany. <https://www.gbif.org/species/1548455>.

**\**Megaselia subcarpalis* (Lundbeck, 1820)**

**Material examined.** DR 14.08. ♂, 11.09. 2 ♂, 26.09. ♂; HR 21.07. 2 ♂, 11.09. ♂, 14.09. 2 ♂♂, 26.09. ♂.

**GBIF records.** There are several records from Sweden and Finland. <https://www.gbif.org/species/1546628>.

**Comments.** The larval diet is unknown (Durska 2013).

**\**Megaselia subconvexa* (Lundbeck, 1920)**

**Material examined.** DR 28.08. ♂, 07.09. ♂; HR 22.08. ♂, 25.08. ♂, 07.09. 2 ♂♂, 11.09. 2 ♂♂.

**Published records.** This species is only known from Denmark and the Netherlands (Schmitz 1941), from Sweden (Bonet et al. 2011), and from Germany (Hannig et al. 2023).

**GBIF records.** There are several records from Sweden, Britain, and Germany. <https://www.gbif.org/species/1547641>.

***Megaselia subtumida* (Wood, 1909)**

**Material examined.** HR 24.07. ♀, 28.07. ♀, 31.07. ♀, 07.08. ♀, 07.09. ♀.

**GBIF records.** In Europe this species is primarily recorded in the northern regions, with some records also from Central Europe. <https://www.gbif.org/species/1547985>.

**\**Megaselia surdifrons* (Wood, 1909)**

**Material examined.** DR 22.09. ♂.

**Published records.** Historical records exist from Denmark, the Netherlands, and Switzerland (Schmitz 1941), while more recent findings have been reported from Norway (Disney 2015) and Germany (Hannig et al. 2023).

**GBIF records.** There are several records from Sweden and Denmark. <https://www.gbif.org/species/1547995>.

**\**Megaselia sylvatica* (Wood, 1910)**

**Material examined.** DR 11.09. ♂, 13.10. ♂.

**GBIF records.** There are several records from Finland, Sweden, and Britain. <https://www.gbif.org/species/1547326>.

**Comments.** The larvae of this species have been reported to be mycophagous (Durska 2013, 2015, 2020) and have been reared from mushrooms belonging to the families Pleurotaceae and Plutaceae (Disney 1991).

***Megaselia vernalis* (Wood, 1909)**

**Material examined.** HR 03.10. ♂.



**GBIF records.** There are several records from Finland, Sweden, Germany, and Britain. <https://www.gbif.org/species/1547053>.

**Comments.** The first Slovak record of this species was published by Grundmann et al. (2023).


## Discussion

This study does not aim to provide a statistically verifiable comparison between the method employed here and commonly used techniques for sampling flying insects. Such a comparison would have required the simultaneous deployment of Malaise traps or yellow pan traps at multiple comparable locations within the zoo. Instead, the primary objective is to demonstrate that the sustainable use of non-target organisms can yield valuable faunistic data. This is particularly relevant for taxa that remain relatively understudied, such as the Phoridae. Nor do we attempt to speculate whether the same species would have been detected using alternative trapping methods. Environmental factors such as vegetation, soil type, microclimate, and the surrounding landscape context exert at least as much influence on species detection as the choice of sampling method itself (Matthews and Matthews 1971). For a comprehensive and detailed analysis of various arthropod sampling techniques, see Schuch et al. (2020).

Depending on the research objective and the methodology applied, investigations involving automated sampling systems typically generate varying amounts of bycatch (non-target organisms) which in most cases remain unanalysed. Only a few projects adopt a holistic approach, aiming to identify and assess as many taxa as possible (Hannig et al. 2023). This study instead seeks to demonstrate that the targeted evaluation of individual, selectively chosen taxa can yield valuable faunistic insights, even in cases where statistical analysis is not feasible. In this study, we present the results obtained using two BG-Sentinel 2 traps combined with CO<sub>2</sub> as an attractant. A total of 73 species of the family Phoridae – represented by nearly 460 individuals – were recorded from these traps between July and October 2023 at Košice Zoo, located in eastern Slovakia. Of these, 24 species are reported for the first time in the Slovak fauna (Fig. 3), increasing the total number of known phorid species in the country from 230 (Mocek 1997, 2009; Jančík and Disney 2020; Grundmann et al. 2023, 2024) to 254.

For comparison, several other studies on Phoridae have reported markedly different results depending on sample size, trapping effort, and methodology. Grundmann and Kappert (2023) identified 71 species (68 from Malaise traps and three from other collection methods) based on a total of nearly 24,000 individuals. Brown and Hartop (2017) recorded 99 species from approximately 42,000 individuals, while Durska (2009) reported 52 species from a sample of 6,000 individuals. In contrast, the present study recorded 73 species from fewer than 500 individuals, highlighting the potential efficiency of the method used. To the best of the authors' knowledge, this is the first application of BG-Sentinel 2 traps with CO<sub>2</sub> for phorid sampling. While this approach may not yield large specimen numbers, it appears to be effective in detecting a high diversity of species, including a significant proportion that are new to the national fauna. Notably, Beuk (2023), using Malaise traps in the Jean Massart Botanic Garden (Auderghem), collected only 126 specimens representing 24 species in 11 genera – further suggesting that even traditional, widely used methods may yield limited results under certain





**Figure 3.** Comparison of the known Slovak phorid fauna (230 species published up to 2024) with the species recorded in the present study (73 species in total, of which 24 are new for Slovakia). The overlapping area represents 49 species, accounting for approximately 21% of the previously documented Slovak phorid fauna.

environmental conditions or in short sampling periods. These results support the idea that greater attention should be paid to non-target organisms collected during entomological surveys, especially when using automated or passive trapping systems. While such material is often regarded as waste, it can represent a significant and largely untapped resource for biodiversity studies, particularly in underexplored or hyperdiverse insect groups such as the Phoridae. We therefore encourage researchers to consider incorporating the analysis of bycatch into their study designs whenever feasible. Even if comprehensive identification of all non-target taxa is not possible, targeted examination of selected groups (such as phorid flies) can yield valuable faunistic, ecological, and biogeographical insights. This approach aligns with the broader goals of sustainability and resource efficiency in biodiversity research, maximising the scientific return from existing sampling efforts. Furthermore, the reuse and re-evaluation of previously overlooked material may facilitate cross-disciplinary collaborations between experts focused on different insect taxa. As demonstrated in this study, what was initially collected for mosquito surveillance turned out to be a rich source of novel data on scuttle flies – revealing both new country records and significant species diversity.

## Conclusions

This study demonstrates that even non-traditional sampling methods, such as BG-Sentinel traps primarily designed for mosquito monitoring, can yield valuable insights into the diversity of non-target insect groups. The discovery of 24 previously unrecorded phorid species for Slovakia emphasises the untapped potential of bycatch material in biodiversity research. This approach not only improves the efficiency of sampling efforts but also opens new avenues for discovering and documenting understudied taxa.

We recommend that future entomological surveys incorporate the analysis of bycatch to enhance the breadth of faunistic data collected. By doing so, researchers can maximise the scientific value of existing collection methods, uncover hidden biodiversity, and foster cross-disciplinary collaboration. Ultimately, the re-evaluation of “unwanted” samples can contribute significantly to our understanding of insect diversity and ecology.



## Acknowledgements

Our thanks go to Jürgen Kappert, for permission to use his beautiful photograph (<http://www.spessart-fliegen.de>). We would especially like to thank the editor and reviewers for providing constructive comments and for improving the manuscript.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This work was co-funded by the European Union under the project 101132974 - OH SUR-Vector. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or European Health and Digital Executive Agency (granting authority). Neither the European Union nor the granting authority can be held responsible for them. This work was supported by the Grant Agency for Doctoral Students and Young Researchers of the University of Prešov (GaPU 11/2024).

### Author contributions

Conceptualisation: BG, JO. Data curation: BG, JO, KLP, TC. Investigation: BG, KLP, TC, LM, PP, JO. Methodology: BG, JO. Validation: BG, KLP, TC, LM, PP, JO. Visualisation: BG. Writing – original draft: OK, JO. Writing – review and editing: BG, KLP, TC, LM, PP, JO.

### Author ORCIDs

Bernd Grundmann  <https://orcid.org/0009-0007-8478-0188>

Katarína Loziaková Peňazziová  <https://orcid.org/0000-0002-3434-5977>

Tomáš Csank  <https://orcid.org/0000-0003-2055-9293>

Laura Mlynárová  <https://orcid.org/0009-0004-1817-7792>

Jozef Oboňa  <https://orcid.org/0000-0002-1185-658X>

### Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Beuk P (2023) Phoridae (Diptera) in the Botanic Garden Jean Massart: new species record for the Belgian fauna and some notes on relationships with ants (Hymenoptera: Formicidae) and ladybirds (Coleoptera: Coccinellidae). In: Grootaert P, Drumont A. (Eds) Flies in the centennial Botanic Garden Jean Massart (Brussels-Capital Region, Belgium). Belgian Journal of Entomology 134: 235–245.
- Bonet J, Ulefors SO, Viklund B, Pape T (2011) Species richness estimations of the megadiverse scuttlefly genus *Megaselia* (Diptera, Phoridae) in a wildfire-affected hemiboreal forest. Insect Science 18(3): 325–348. <https://doi.org/10.1111/j.1744-7917.2010.01362.x>



- Brown BV, Hartop EA (2017) Big data from tiny flies: patterns revealed from over 42,000 phorid flies (Insecta: Diptera: Phoridae) collected over one year in Los Angeles, California, USA. *Urban Ecosystems* 20(3): 521–534. <https://doi.org/10.1007/s11252-016-0612-7>
- Brown BV, Marshall SA (1984) Phorid flies (Diptera: Phoridae) associated with mushrooms in Southern Ontario. *Proceedings of the Entomological Society of Ontario* 115: 77–80.
- Büchs W (1988) Stamm- und Rinden-Zoozönosen verschiedener Baumarten des Hartholzauenwaldes und ihr Indikatorwert für die Früherkennung von Baumschäden. Ph. D. Thesis Rheinische Friedrich-Wilhelms-Universität Bonn, 813 pp.
- Buck M, Disney RHL (2001) Revision of the *Megaselia giraudii* and *M. densior* species complexes of Europe, including ecological notes (Diptera, Phoridae). *Beiträge zur Entomologie* 51(1): 73–154. <https://doi.org/10.21248/contrib.entomol.51.1.73-154>
- Caruso V, Hartop E, Chimeno C, Noori S, Srivathsan A, Haas M, Lee L, Meier R, Whitmore D (2024) An integrative framework for dark taxa biodiversity assessment at scale: A case study using *Megaselia* (Diptera, Phoridae). *Insect Conservation and Diversity* 17(6): 968–987. <https://doi.org/10.1111/icad.12762>
- Disney RHL (1983) Scuttle flies – Diptera, Phoridae (except *Megaselia*). *Handbooks for the Identification of British Insects* 10(6): 1–81.
- Disney RHL (1989) Scuttle Flies – Diptera Phoridae Genus *Megaselia*. *Handbooks for the Identification of British Insects* 10(8): 1–155.
- Disney RHL (1991) Phoridae. In: Soos A, Papp L (Eds) *Catalogue of Palaearctic Diptera*. Vol. 7. Dolichopodidae – Platypezidae – Akademiai Kiado, Budapest, 143–204. <https://doi.org/10.1016/B978-0-444-98731-0.50010-7>
- Disney RHL (1994) *Scuttle flies: the Phoridae*. Chapman, Hall, London, Glasgow, 467 pp. <https://doi.org/10.1007/978-94-011-1288-8>
- Disney RHL (1999) A troublesome sibling species complex of scuttle flies (Diptera, Phoridae) revisited. *Journal of Natural History* 33(8): 1159–1216. <https://doi.org/10.1080/002229399299987>
- Disney RHL (2004) The distinctive *Megaselia deltomera* Schmitz, 1924 (Diptera: Phoridae) belongs to a complex of sibling species. *Annales Zoologici* 54(4): 687–695.
- Disney RHL (2006a) Revision of the Palaearctic members of the species complex resembling *Megaselia brevior* (Diptera: Phoridae). *Fragmenta Faunistica* 49(1): 41–51. <https://doi.org/10.3161/00159301FF2006.49.1.041>
- Disney RHL (2006b) A new species of *Megaselia* Rondani (Dipt., Phoridae) from Britain and a new synonym. *Entomologist's Monthly Magazine* 142: 31–39. <https://doi.org/10.31184/M00138908.1584.4154>
- Disney RHL (2010a) Scuttle flies (Diptera: Phoridae) reared from logs in Finland and N.W. Russia, including two new species. *Entomologica Fennica* 20(4): 257–267. <https://doi.org/10.33338/ef.84487>
- Disney RHL (2010b) Recognition of four poorly known species of *Megaselia* Rondani (Dipt., Phoridae). *Entomologist's Monthly Magazine* 146: 143–147.
- Disney RHL (2015) Scuttle flies (Diptera, Phoridae) from the canopies of oak trees (Fagaceae) in Norway, including 13 new species. *Norwegian Journal of Entomology* 62: 20–52. <https://doi.org/10.13102/sociobiology.v62i1.124-127>
- Disney RHL, Evans RE (1982) Records for Phoridae (Diptera) reared from fungi. *Entomologist's Record and Journal of Variation* 94: 104–105.
- Disney RHL, Withers P (2011) Scuttle flies (Diptera, Phoridae) reared from tree rotholes in France, including three new species of *Megaselia* Rondani. *Fragmenta Faunistica* 54(1): 29–41. <https://doi.org/10.3161/00159301FF2011.54.1.029>



- Disney RHL, Coulson JC, Butterfield J (1981) A survey of the scuttle flies (Diptera: Phoridae) of upland habitats in northern England. *Naturalist* 106: 53–66.
- Durska E (2009) The scuttle fly (Diptera: Phoridae) assemblages of pine plantations of the Biała Forest (Poland). *Entomologica Fennica* 20(3): 170–178. <https://doi.org/10.33338/ef.84473>
- Durska E (2013) Effects of disturbances on scuttle flies (Diptera: Phoridae) in pine forests. *Biodiversity and Conservation* 22(9): 1991–2021. <https://doi.org/10.1007/s10531-013-0522-4>
- Durska E (2015) Effects of fire on scuttle flies (Diptera: Phoridae) in a pine forest in Poland. *Entomologica Fennica* 26(4): 181–193. <https://doi.org/10.33338/ef.84645>
- Durska E (2020) Preliminary data of the scuttle flies (Diptera: Phoridae) in the linden-oak-hornbeam forest of the Wigry National Park, North-East-Poland. *Fragmenta Faunistica* 63(1): 47–52. <https://doi.org/10.3161/00159301FF2020.63.1.047>
- Grundmann B, Kappert J (2023) The Phoridae (Diptera) of NE-Westphalia: A field study over five years. *Fragmenta Faunistica* 66(1): 15–36. <https://doi.org/10.3161/00159301FF2023.66.1.015>
- Grundmann B, Oboňa J, Mocek B, Mock A (2023) Scuttle flies (Diptera: Phoridae) of subterranean habitats in Slovakia: a review and a case study. *Biologia* 78(12): 3591–3597. <https://doi.org/10.1007/s11756-023-01486-4>
- Grundmann B, Manko P, Oboňa J (2024) Overlooked insects in neglected ecosystem: New records of Phoridae for Slovakia discovered in rural environment. *Historia Naturalis Bulgarica* 46(4): 109–118. <https://doi.org/10.48027/hnb.46.043>
- Hannig K, Ehlert T, Fuhrmann M, Grundmann B (2023) Zur Fauna und Flora einer Sandabgrabung bei Haltern-Flaesheim (Kreis Recklinghausen, Nordrhein-Westfalen) – 3. Nachtrag. In: Hannig K, Kriegs JO, Tenbergen B (Eds) Einige Besonderheiten und Dynamik der Flora und Fauna Nordrhein-Westfalens. *Abhandlungen aus dem Westfälischen Museum für Naturkunde* 106: 7–162.
- Jančík L, Disney RHL (2020) A new species of *Aenigmatias* Meinert (Diptera: Phoridae) from Slovakia, whose larvae parasitise a queen ant (Hymenoptera: Formicidae). *Entomologist's Monthly Magazine* 156(2): 103–108. <https://doi.org/10.31184/M00138908.1562.3965>
- Langourov M (2004a) Scuttle flies (Diptera: Phoridae) from urban and suburban areas in the Sofia Plain. In: Penev L, Niemelä J, Kotze DJ, Chipev N (Eds) *Ecology of the City of Sofia. Species and Communities in an Urban Environment*, Pensoft Publishers, Sofia-Moscow, 429–436.
- Langourov M (2004b) Scuttle flies (Diptera: Phoridae) from Biogradska Gora National Park-Montenegro. *The Biodiversity of the Biogradska Gora National Park. Monographies I*, Department of Biology, University of Montenegro and Centre for Biodiversity of Montenegro, 104–112.
- Matthews RW, Matthews JR (1971) The Malaise Trap: Its Utility and Potential for Sampling Insect Populations. *Great Lakes Entomologist* 4: 117–122. <https://doi.org/10.22543/0090-0222.1158>
- Mocek B (1997) Faunistic records from the Czech and Slovak Republics: Diptera, Phoridae. In: Vaňhara M, Rozkošný V (Eds) *Dipterologica Bohemoslovaca*, 8. *Folia Fac Sci Nat Univ Masarykianae Brunensis, Biologia* 95: 220–222.
- Mocek B (2009) Phoridae Latreille, 1796. In: Jedlička L, Kúdela M, Stloukalová V (Eds) *Checklist of Diptera of the Czech Republic and Slovakia*. <http://www.edvis.sk/diptera2009/families/phoridae.htm> [accessed 31.10.2023]



- Oosterbroek P (2006) The European Families of the Diptera: Identification-Diagnosis-Biology. KNNV Publishing, Utrecht, 205 pp. [https://doi.org/10.1163/9789004278066\\_013](https://doi.org/10.1163/9789004278066_013)
- Peňazziová K, Korytár L, Pastorek P, Pistl J, Rusňáková D, Szemes T, Čabanová V, Ličková M, Boršová K, Klempa B, Csank T (2021) Genetic Characterization of a Neurovirulent West Nile Virus Variant Associated with a Fatal Great Grey Owl Infection. *Viruses* 13(4): 699. <https://doi.org/10.3390/v13040699>
- Prescher S, Bellstedt R (1994) Beitrag zur Kenntnis der Buckelfliegenfauna Thüringens (Dipt., Phoridae). *Entomologische Nachrichten und Berichte* 38: 45–51.
- Schmitz H (1941) Kritisches Verzeichnis der Paläarktischen Phoriden mit Angabe ihrer Verbreitung (Fortsetzung). *Natuurhistorisch Maandblad* 30: 11–12; 15–17; 34–35.
- Schmitz H (1943) 33. Phoridae (1. Teilband). Lieferung 147+149. In: Lindner E (Ed.) *Die Fliegen der paläarktischen Region* 4(33): 129–192.
- Schmitz H (1951) 33. Phoridae (1. Teilband). Lieferung 165. In: Lindner E (Ed.) *Die Fliegen der paläarktischen Region* 4(33): 241–272.
- Schmitz H (1956) 33. Phoridae (1. Teilband). Lieferung 187. In: Lindner E (Ed.) *Die Fliegen der paläarktischen Region* 4(33): 369–416.
- Schuch S, Ludwig H, Wesche K (2020) Erfassungsmethoden für ein Insektenmonitoring. Eine Materialsammlung. Bundesamt für Naturschutz, BfN-Schriften 565: 85.
- Szadziewska MM (1977) Geofilne muchówki (Diptera) rezerwatu 'Las Piwnicki' koło Torunia. *Acta Universitatis Nicolai Copernici Biologia* 19(39): 107–115.
- Weber G, Prescher S, Ulefors S-O, Viklund B (2006) Fifty-eight species of scuttle flies (Diptera: Phoridae: *Megaselia* spp.) new to Sweden from the (Tyrestra) Tyresta National Park and Nature Reserve. *Studia Dipterologica* 13: 231–240.